



Agenda		
4:30 – 4:45 pm	Challenge of Adherence in Pharma:	: Case Study
4:45 – 5:30 pm	Breakout Groups – Applying the Evidence	
5:30 – 6:00 pm	Team Presentation & Discussion	
FOCUS	©2006 - 2016 Return on Focus LLC. All rights reserved. PROPRIETARY AND CONFIDENTIAL 2	RUTCEERS Interdisciplinary Job Opportunities for Biomedical Scientists (UO85)





Patient Adherence Rates Haven't Changed **Substantially** 

"Adherence rates average around 50% and range from 0% to over 100%, and there is no evidence for substantial change in the past 50 years."

-Nieuwlaat et al. 2014

Source: Niewwlaat et al, Interventions for Enhancing Medication Adherence - Cochrane Collaboration, 2014.

## **Definition of Adherence**

Adherence:


The extent to which the patient continues the agreed-upon mode of treatment under limited supervision when faced with conflicting demands, as distinguished from compliance or maintenance.


Source: The American Heritage Medical Dictionary online at www.medical.yourdictionary.com/adherence. Accessed December 2015.


RETURN ON FOCUS


©2006 - 2016 Return on Focus LLC. All rights reserved. PROPRIETARY AND CONFIDENTIAL 6













## Physicians Key Culprit to Adherence Throughout the Treatment Continuum



Physician impact on non-adherence is multi-factorial and is often substantially underestimated


	Treatment Initiation	
MDs incorrectly believe the can identify which patients will be adherent and which will not, and they make prescribing decisions based on these assumptions <sup>1</sup>	MDs do not typically	Treatment Continuation
	communicate the criticality of adherence <sup>1</sup> Frequently tell patients occasionally missing doses is not a significant problem	MDs are reluctant to mention adherence during consults for fear of damaging patient trust <sup>2</sup> Do not always reduce dosage to manage side effects or increase dosage to improve efficacy <sup>3</sup>
Medical Research and Opinion, 2009. 2. D	enois VR, et al. "Adherence with oral chemo	to adhere to prescription medications for chi therapy: results from a qualitative study of th t al. Adherence to Medication. <i>NEJM</i> , August







## **Gleevec Case Study** You're the Brand Manager for Gleevec for the treatment of CML (chronic myeloid leukemia) • Before the introduction of Gleevec the survival rate of CML patients after one Gleevec" year was between 25% to 50% 100 Gleevec changed CML from a fatal cancer into a manageable chronic condition • 5 Year survival rate is 95% "You would think that cancer patients would be more motivated to keep taking their drug, and so Non-adherence to Gleevec therapy the finding [non-adherence] is rather affects cytogenic and molecular counterintuitive" - Dr. Nick Barber responses CML patients are no more adherence to their medication than patients on high blood pressure medications FOCUS ©2006 - 2016 Return on Focus LLC. All rights reserved. PROPRIETARY AND CONFIDENTIAL isciplinary Job Opportunitio medical Scientists (IJOBS) 14


## **Gleevec Adherence Rates Range from 44% to 79%**








Published Adherence Rates for Gleevec\*


























©2006 - 2016 Return on Focus LLC. All rights reserved. PROPRIETARY AND CONFIDENTIAL 26

